

AD-A131 624

COLLECTION SEPARATION CRYOPRESERVATION AND
CHARACTERIZATION OF PERIPHERAL (U) BOSTON UNIV MA
SCHOOL OF MEDICINE A J ROY ET AL. 01 OCT 82


1/1


UNCLASSIFIED

N00014-79-C-0489

F/G 6/5

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

ADA 131624

"A"
To DTIC

OFFICE OF NAVAL RESEARCH

Contract N00014-7⁹-C-0489

Task No. NR207-196

ANNUAL REPORT

June 1, 1981 to May 31, 1982

COLLECTION, SEPARATION, CRYOPRESERVATION AND CHARACTERIZATION
OF PERIPHERAL BLOOD AND BONE MARROW STEM CELLS AND
THEIR USE IN TREATING LETHALLY IRRADIATED DOGS

by

Albert J. Roy, Ph.D.
[REDACTED] Research Laboratory
Boston University School of Medicine
615 Albany Street
Boston, MA 02118

1 October, 1982

Reproduction in whole or in part is permitted for any purpose of the
United States Government

Distribution of this report is unlimited

DTIC
SELECTED
NOV 1983
D
E

THE BETH H. MOORE
LIBRARY


DTIC FILE COPY

88 02

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Annual Report No. 1	2. GOVT ACCESSION NO. AD A131624	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Cryopreservation of Canine Peripheral Blood Mononuclear Cells in Untreated and Ficoll-Hypaque Treated Buffy Coats		5. TYPE OF REPORT & PERIOD COVERED 6/1/81-5/31/82
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) A.J. Roy, A.J. Melaragno, L.T. Lavigne, Jr., A.D. Gray, J. Dittmer, and C.R. Valeri		8. CONTRACT OR GRANT NUMBER(s) 9 ONR N00014-78-C-0489
9. PERFORMING ORGANIZATION NAME AND ADDRESS Boston University School of Medicine Naval Blood Research Laboratory Boston, MA. 02118		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR207-196 Code 444
11. CONTROLLING OFFICE NAME AND ADDRESS Biological Sciences Division Office of Naval Research (N00014) 800 N. Quincy Street, Arlington, Va. 22217		12. REPORT DATE 1 October, 1982
		13. NUMBER OF PAGES 20
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Same as above		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE NA
16. DISTRIBUTION STATEMENT (of this Report) Distribution of this report is unlimited and is not to be distributed outside the Department of Defense		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) Same as above		
18. SUPPLEMENTARY NOTES NA		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) blood mononuclear cells, cryopreservation, mononuclear cell harvest, ficoll-hypaque separation, bone marrow repopulation		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Peripheral blood mononuclear cells (MNC) were obtained from buffy coats collected during 21 platelet apheresis procedures in the dog using the Haemonetics 30 Blood Processor. The mononuclear cells in the buffy coat were divided into two equal portions. One portion was treated with ficoll-hypaque to purify the mononuclear cells by removing the granulocytes and red cells. The other portion was not treated. Dimethylsulfoxide (Me ₂ SO) in McCoy's medium was added to the untreated buffy coats and to the ficoll-		

hypaque treated buffy coats rapidly in 1 to 2 minutes or slowly over 15 to 20 minutes. The cell suspensions were frozen in polyolefin plastic bags at 2-3 C per minute by placing the plastic bag in a -80 C freezer, or at 1 C per minute by use of a graded freezing apparatus. The percentage of viable MNC's was determined after thawing and washing by measurement of uptake of fluorescein diacetate and ethidium bromide. There was no significant difference in the percentage of viable mononuclear cells recovered between the untreated buffy coat and the ficoll-hypaque treated buffy coat depleted of contaminating red blood cells and granulocytes regardless of the rate of addition of the Me₂SO-McCoy's solution and the rate of freezing. There was a significant difference in the number of viable MNC's recovered when comparing untreated buffy coat and ficoll-hypaque treated buffy coat due to a 22% loss of MNC's during ficoll-hypaque treatment.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Avail and/or	
Dist	Special
A	



CRYOPRESERVATION OF CANINE PERIPHERAL BLOOD MONONUCLEAR CELLS
IN UNTREATED AND FICOLL-HYPAQUE TREATED BUFFY COATS

A.J. ROY, A.J. MELARAGNO, L.T. LAVIGNE, JR, A.D. GRAY, J. DITTMER
AND C.R. VALERI

NAVAL BLOOD RESEARCH LABORATORY AND
DEPARTMENTS OF MEDICINE AND ANATOMY
BOSTON UNIVERSITY SCHOOL OF MEDICINE
BOSTON, MA, 02118

Prepared for Publication
in
Vox Sanguinis

RUNNING TITLE: CRYOPRESERVATION OF CANINE BLOOD MONONUCLEAR CELLS

KEY WORDS:

Blood mononuclear cells
Cryopreservation
mononuclear cell harvest
Ficoll-hypaque separation
Bone marrow repopulation

Please address correspondence to:

Dr. Albert J. Roy
Naval Blood Research Laboratory
Boston University School of Medicine
615 Albany Street
Boston, MA, 02118
(617-247-6700)

SUMMARY

Peripheral blood mononuclear cells (MNC) were obtained from the buffy coats collected during 21 platelet apheresis procedures using the Haemonetics 30 Blood Processor. The mononuclear cells in the buffy coat were divided into two equal portions. One portion was treated with ficoll-hypaque to purify the mononuclear cells by removing the granulocytes and red cells, the other portion was not treated. Dimethylsulfoxide (Me_2SO) in McCoy's medium was added to the untreated buffy coats and to the ficoll-hypaque treated buffy coats rapidly in 1 to 2 minutes or slowly over 15 to 20 minutes. The cell suspensions were frozen in polyolefin plastic bags at 2-3 C/ min by placing the plastic bag in a -80 C freezer, or at 1 C/min by use of a graded freezing apparatus. The percentage of viable MNC's was determined after thawing and washing by measurement of uptake of fluorescein diacetate and ethidium bromide.

There was no significant difference in the percentage of viable mononuclear cells recovered between the untreated buffy coat and the ficoll-hypaque treated buffy coat depleted of contaminating red blood cells and granulocytes regardless of the rate of addition of the Me_2SO -McCoy's solution and the rate of freezing. There was a significant difference in the number of viable MNC's recovered when comparing untreated buffy coat and ficoll-hypaque treated buffy coat due to a 22% loss of MNC's during ficoll-hypaque treatment.

INTRODUCTION

The transfusion of mononuclear cells obtained from the peripheral blood rather than from the bone marrow to repopulate marrow damaged by ionizing radiation or by chemicals is currently being studied.⁽¹⁻³⁾ We have recently demonstrated that the mononuclear cells in buffy coat can be isolated during platelet apheresis procedures in the dog using the Haemonetics 30 discontinuous flow centrifugal method.⁽⁴⁾ Autologous transfusion of bone marrow or peripheral blood MNC's has been recommended in patients when compatible donors are not available.⁽⁵⁾ Mononuclear cells obtained from peripheral blood or bone marrow contaminated with red blood cells and granulocytes have been administered as autologous transfusions. Investigators have recommended the removal of red blood cells and granulocytes from mononuclear cells obtained from peripheral blood and bone marrow by treatment with ficoll-hypaque gradient centrifugation⁽⁶⁾ The limited number of totipotent stem cells in peripheral blood necessitates several apheresis procedures to isolate enough mononuclear cells to successfully repopulate the bone marrow.⁽⁴⁾ These totipotent hematopoietic peripheral blood mononuclear cells must be frozen as they are collected in order to obtain an adequate number for subsequent transfusion.

The present study was designed to determine the loss of peripheral blood mononuclear cells during treatment with ficoll-hypaque and the recovery of viable mononuclear cells in untreated buffy coats and ficoll-hypaque treated buffy coats processed by the rapid or slow addition of Me₂SO-McCoy's cryopreservative solution and by freezing at 1C/min to -40C or at 2-3C/min to -80C in polyolefin plastic bags prior to storage at -150C followed by thawing and washing.

MATERIALS AND METHODS

Collection of Buffy Coat During Discontinuous Flow Centrifugation

Healthy beagle dogs weighing 10 to 15 kg were studied. At least two weeks before their first apheresis procedure, an arterio-venous fistula was surgically created between a carotid artery and jugular vein to allow adequate blood flow into the Haemonetics 30 Blood Processor. A 16 gauge catheter was inserted into the venous end of the fistula in anesthetized dogs and blood was collected into a 125 ml pediatric bowl. Blood was delivered into the spinning bowl at 60 ml/min. and mononuclear cell collection was begun when the top of the red cell volume was $1\frac{1}{2}$ to 2 cm from the core of the bowl. the blood flow was then reduced to 20 ml/min. Mononuclear cell collection continued for 45 seconds after red blood cells appeared in the collection port. Plasma and red blood cells were returned to the dog through an 18 gauge catheter inserted into a vein in the foreleg. This process was repeated ~~three~~ times for a total of four "passes". Acid-citrate-dextrose (Formula A) anticoagulant was used in a ratio of 1 volume of anticoagulant to 7 volumes of blood.

Processing of the Buffy Coat

The buffy coat containing platelets, mononuclear cells, granulocytes and red cells was centrifuged at $160 \times g$ for 10 minutes in a Sorval RC3B centrifuge maintained at $22 \pm 2^\circ C$. The platelet rich plasma (PRP) depleted of red blood cells and leukocytes was expressed into a 600 ml polyvinyl-chloride transfer pack. The platelet count was measured in duplicate using phase-contrast microscopy and the platelets in the PRP were frozen as described previously.⁽⁷⁾ An aliquot of the residual buffy coat was obtained.

White blood cell counts were measured in triplicate using the Coulter model Zf. Smears were prepared in duplicate for differential white blood cell counts. The number of mononuclear cells was determined by multiplying the white blood cell count by the percentage of mononuclear cells on the smears and by the volume of the residual buffy coat. This buffy coat was divided into two equal aliquots. One of these was frozen as described below without further treatment. The second was treated by ficoll-hypaque density centrifugation to remove the granulocytes and red cells. This sample was diluted with Hank's balanced salt solution (HBBS) to a concentration of approximately 2×10^7 leukocytes. A 5 ml volume of diluted residual buffy coat was layered over a 3 ml volume of a ficoll-hypaque solution with a density of 1.077 in 15 ml conical centrifuge tubes. The sample was centrifuged at 1500 rpm ($500 \times g$) for 40 minutes at room temperature in a Damon CRU-5000 centrifuge using a #269 head. The mononuclear cell layers were collected into 50 ml conical centrifuge tubes and diluted with HBBS. After mixing well, the samples were again centrifuged at 1500 rpm for 10 minutes. The supernatant was decanted and the pellets were resuspended in HBBS. This process was repeated again and the cells were then resuspended in McCoy's medium to a volume of 20 ml. An 0.3 ml sample was taken for total WBC count and to prepare smears for differential white blood cell counts. The total number of mononuclear cells was calculated as described above.

Cryopreservation of MNC's

A volume of 20% Me₂SO in McCoy's medium equal to the volume of untreated buffy coat or ficoll-hypaque treated buffy coat was prepared as follows: Five volumes of McCoy's medium kept in wet ice at $\pm 4^\circ \text{C}$ were pipetted into a sterile bottle which was also kept in ice. One volume of Me₂SO was added to the McCoy's medium by syringe in about one minute with constant manual spinning of the bottle to dissipate the heat generated. A volume of the 20% Me₂SO-McCoy's

mixture was added to an equal volume of the cell suspension, transferred aseptically to a polyolefin freezing bag through a sampling site coupler (Fenwal #4C2405) to achieve a final Me_2SO concentration of 10%. Addition of the cryoprotectant solution was accomplished in one of two ways:

- (a) rapid addition in 1 to 2 minutes with constant agitation of the bag on ice.
- (b) slow addition over 15 to 20 minutes with constant agitation of the bag on ice; mixing was accomplished using an Eberbach shaker adjusted to run at 180 lateral oscillations per minute.

The volume of buffy coat-cryoprotectant was approximately 80 ml; after isolation of the ficoll-hypaque separated buffy coat, the cell-cryoprotectant volume was about 40 ml.

The polyolefin plastic bag was placed in an aluminum freezing container and frozen in one of two ways:

- (a) a freezing rate of 2-3 $^{\circ}\text{C}/\text{min}$ by placement into a -80°C mechanical freezer for 12 hours followed by transfer of the container into the gas phase of a liquid nitrogen refrigerator maintained at -150°C .
- (b) a freezing rate of 1 $^{\circ}\text{C}/\text{min}$ by freezing in a Cryo-Med graded freezing apparatus from $+4$ to -40°C followed by transfer of the container into the gas phase of a liquid nitrogen refrigerator maintained at -150°C .

Three freezing combinations were evaluated in this study:

- (a) rapid addition of cryoprotectant and freezing at 2-3 $^{\circ}\text{C}/\text{min}$.
- (b) slow addition of cryoprotectant and freezing at 2-3 $^{\circ}\text{C}/\text{min}$.
- (c) slow addition of cryoprotectant and freezing at 1 $^{\circ}\text{C}/\text{min}$.

Thawing and Washing Procedure

The concentrates were thawed by immersion of the bag in a 37°C water bath with constant manual agitation until the last ice particle had melted. This was accomplished in less than 30 seconds. An 0.5 ml sample was obtained

aseptically by syringe and needle for white blood cell and differential counts and for viability testing. Fifty ml of McCoy's medium was added rapidly by needle through a sampling port. The diluted cell suspension was transferred into a 300 ml capacity polyvinylchloride bag and centrifuged at 4160 rpm (4470 x g) for 5 minutes at room temperature in a Sorval RC-33 centrifuge. The supernatant was removed using a sterile Fern #4C2244 transfer set and 35 to 50 ml of platelet poor plasma was added. The cells were resuspended by gentle manual manipulation of the bag. A second 0.5 ml sample was obtained for testing.

The tests performed after thawing and washing were as follows:

- (a) total WBC count in triplicate using the Coulter model Zf counter.
- (b) differential WBC counts in duplicate on slides stained with Wright's-Giemsa.
- (c) viability was assessed with fluorescein diacetate and ethidium bromide, as described previously. (8,9)

RESULTS

The percentage recovery of viable peripheral blood mononuclear cells in untreated buffy coat after rapid addition of the cryoprotective agent and freezing at 2-3 C/min, thawing and washing was 74%. (Table 1) The percentage recovery of viable peripheral blood mononuclear cells in ficoll-hypaque treated buffy coat frozen in an identical manner was 70%. (Table 1) When the Me₂SO-McCoy's medium was added slowly to the cell suspension and freezing was carried out at 2-3 C/min the recovery of viable MNC's in untreated buffy coat averaged 54% and ficoll-hypaque treated buffy coats averaged 60%. (Table 2) Freezing at 1C/min following slow addition of the cryoprotective solution resulted in a recovery of 73% viable mononuclear cells with both untreated buffy coat and ficoll-hypaque treated buffy coat. (Table 3) No significant

differences in the percentage recovery of mononuclear cells that were frozen were observed among the three groups. ($p > 0.5$, > 0.2 and > 0.8 respectively by the Student non-paired t test).

Twenty-two percent of the mononuclear cells were lost during the treatment of the buffy coat by ficoll-hypaque density centrifugation to remove granulocytes and red blood cells. (Table 4). The recovery of viable mononuclear cells calculated from the number of cells collected prior to treatment with ficoll-hypaque was 52% when they were frozen at 2-3 C/min after rapid addition of the cryophylactic medium; 36% when frozen at 2-3 C/min after slow addition of the cryophylactic medium; and 57% when frozen at 1 C/min after slow addition of the cryophylactic medium. (Tables 1,2,3).

The percentage recovery of viable mononuclear cells that were collected showed that there were significant differences between non-treated buffy coats and ficoll-hypaque treated buffy coat ($p < 0.05$, < 0.01 , and < 0.05 respectively by the Student non-paired t test) regardless of the rate of addition of the Me_2SO and the rate of freezing. (Tables 1,2,3).

DISCUSSION

Recovery of more than 70% viable peripheral blood dog mononuclear cells in non-treated buffy coat was observed in our previous study whether freezing with 10% Me_2SO was achieved at 1 C/min using a graded freezing apparatus or at 2-3 C/min by storage in a -80 C mechanical freezer. (4)

In the studies reported here, when the percentage recovery of viable mononuclear cells was related to the number frozen, the freeze-thaw-wash recovery of viable peripheral blood mononuclear cells was similar for the untreated and the ficoll-hypaque treated buffy coat and was independent of the rate of addition of Me_2SO and the rate of freezing. There was no signif-

icant difference between freezing at 2-3 C/min with rapid addition of the cryophylactic agent as compared to graded freezing at 1 C/min with slow addition of the cryophylactic agent. With both untreated buffy coat and ficoll-hypaque treated buffy coat, however, slow addition of the Me_2SO medium and freezing at 2-3 C/min resulted in a significantly greater loss of viability of mononuclear cells than was seen using the other two freezing processes. The treatment of buffy coat with ficoll-hypaque to remove red cells and granulocytes prior to freezing removed approximately 20% of the mononuclear cells.

The number of viable mononuclear cells following thawing was consistently less than that seen after thawing and washing, whether untreated buffy coat or ficoll-hypaque treated buffy coat was cryopreserved. The increase in the number of viable mononuclear cells following washing was due to an increase in viability as measured by fluorescein diacetate uptake and ethidium bromide exclusion rather than by an increase in recovery of mononuclear cells. The 10% concentration of dimethylsulfoxide in the thawed cell preparations may increase the permeability of the cells to ethidium bromide causing a greater number of cells to be called non-viable. The reduction of the Me_2SO during washing may return the permeability of the mononuclear cells towards normal.

TABLE 1

RECOVERY (MEAN + S.D.) OF PERIPHERAL BLOOD MONONUCLEAR CELLS AFTER
RAPID ADDITION OF A CRYOPHYLACTIC AGENT, FREEZING AT 2-3 C PER MINUTE
THAWING AND WASHING IN 8 DOGS

Source of Mononuclear Cells	Untreated Buffy Coat	Ficoll-Hypaque Treated Buffy Coat
Mononuclear Cells Isolated ($\times 10^9$)	.61 \pm .24	.61 \pm .24
Mononuclear Cells Frozen ($\times 10^9$)	.61 \pm .24	.46 \pm .15*
Number of Mononuclear Cells Viable Post-thaw ($\times 10^9$)	.34 \pm .11	.17 \pm .04
Recovery of Viable Mononuclear Cells After Thawing (%)	56 \pm 22	38 \pm 8
Number of Mononuclear Cells Viable Post-wash ($\times 10^9$)	.45 \pm .22	.32 \pm .09
Recovery of Viable Mononuclear Cells Post-wash (%)		
of the Number Frozen	74 \pm 20	70 \pm 15 ($p > 0.5$)
of the Number Collected	74 \pm 20	52 \pm 14 ($p < .05$)

*The number remaining after ficoll-hypaque treatment

TABLE 2

RECOVERY (MEAN + S.D.) OF PERIPHERAL BLOOD MONONUCLEAR CELLS AFTER
SLOW ADDITION OF A CRYOPHYLACTIC AGENT, FREEZING AT 2-3 C PER MINUTE
THAWING AND WASHING IN 6 DOGS

Source of Mononuclear Cells	Untreated Buffy Coat	Ficoll-Hypaque Treated Buffy Coat
Mononuclear Cells Isolated (x 10 ⁹)	.69 ± .14	.69 ± .14
Mononuclear Cells Frozen (x 10 ⁹)	.69 ± .14	.42 ± .17*
Number of Mononuclear Cells Viable Post-thaw (x 10 ⁹)	.31 ± .06	.23 ± .09
Recovery of Viable Mononuclear Cells After Thawing (%)	46 ± 11	57 ± 11
Number of Mononuclear Cells Viable Post-wash (x 10 ⁹)	.37 ± .11	.25 ± .11
Recovery of Viable Mononuclear Cells Post-wash (%)		
of the Number Frozen	54 ± 13	60 ± 9 (p > 0.2)
of the Number Collected	54 ± 13	36 ± 4 (p < .01)

*The number remaining after ficoll-hypaque treatment

TABLE 3

RECOVERY (MEAN + S.D.) OF PERIPHERAL BLOOD MONONUCLEAR CELLS AFTER
SLOW ADDITION OF A CRYOPHYLACTIC AGENT, FREEZING AT 1 C PER MINUTE
THAWING AND WASHING IN 7 DOGS

Source of Mononuclear Cells	Untreated Buffy Coat	Ficoll-Hypaque Treated Buffy Coat
Mononuclear Cells Isolated ($\times 10^9$)	.70 \pm .08	.70 \pm .08
Mononuclear Cells Frozen ($\times 10^9$)	.70 \pm .08	.55 \pm .08*
Number of Mononuclear Cells Viable Post-thaw ($\times 10^9$)	.34 \pm .08	.32 \pm .08
Recovery of Viable Mononuclear Cells After Thawing (%)	48 \pm 22	53 \pm 22
Number of Mononuclear Cells Viable Post-wash ($\times 10^9$)	.51 \pm .20	.40 \pm .11
Recovery of Viable Mononuclear Cells Post-wash (%)		
of the Number Frozen	73 \pm 16	73 \pm 12 (p > 0.8)
of the Number Collected	73 \pm 16	57 \pm 7 (p < .05)
The number remaining after ficoll-hypaque treatment		

TABLE 4

RECOVERY OF DOG PERIPHERAL BLOOD MONONUCLEAR CELLS FROM 19 BUFFY COATS
TREATED WITH FICOLL-HYPAQUE

	<u>Number of Mononuclear Cells Collected ($\times 10^9$)</u>	<u>Number of Mononuclear Cells Remaining After Treatment ($\times 10^9$)</u>	<u>% Mononuclear Cells Remaining After Treatment</u>
Mean	0.66	0.51	78
Standard Deviation	0.13	0.12	10
Range	0.40 - 1.19	0.28 - 0.78	62 - 100

REFERENCES

1. Goldman, J., Catovsky, D., Goolden, A., Johnson, S. and Galton, D.:
Buffy coat autografts for patients with chronic granulocytic leukemia
in transformation. *Blut*, 42:149-155, (1981)
2. McGann, L., Turner, A., Allalunis, M., and Turc, J.: Cryopreservation
of human peripheral blood stem cells: Optimal cooling and warming con-
ditions. *Cryobiology*, 18:469-472, (1981)
3. Sarpel, S., Zander, A., Harvath, L., and Epstein, R.: The collection,
preservation and function of peripheral blood hematopoietic cells in
dogs. *Exp. Hemat.*, 7:113-120, (1979)
4. Roy, A., Melaragno, A., Lavigne, L., Jr., Gray, A., Dittmer, J. and
Valeri, C.R.: The cryopreservation of canine peripheral blood mono-
nuclear cells obtained as a by-product of platelet collection. In
Preparation
5. Diesseroth, A., and Abrams, R.: The role of autologous stem cell recon-
stitution in intensive therapy for resistant neoplasms. *Cancer Treat.*
Rep., 63:461-471. (1974)
6. Thorsby, E. and Bratlie, A.: A rapid method for the preparation of pure
lymphocyte suspensions, *Histocompatibility Testing*, 1970, ed., P.I.
Terasaki, p 655, Munksgaard, Copenhagen (1970)
7. Melaragno, A., Abdu, W., Katchis, R., Vecchione, J. and Valeri, C.R.:
Cryopreservation of platelets isolated with the IBM 2997 blood cell
separator: a rapid and simplified approach. *Vox Sang.* (In Press)

8. Persidsky, M. and Ballie, G.: Fluorometric test of cell membrane integrity. *Cryobiology*, 14:322-331, (1977)
9. Lionetti, F., Hunt, S., Mattaliano, R. and Valeri, C.R.: In vitro studies of cryopreserved baboon granulocytes. *Transfusion*, 19:685-692, (1978)

ACKNOWLEDGEMENTS

This research was supported by Contracts #N00014-79-C-0489 and
#N000014-C-0168 from the Office of Naval Research.

19

OFFICE OF NAVAL RESEARCH
BIOLOGICAL SCIENCES DIVISION
BIOPHYSICS PROGRAM, Code 444
DISTRIBUTION LIST FOR TECHNICAL, ANNUAL AND FINAL REPORTS

Number of Copies

- (13) Administrator, Defense Documentation Center
Cameron Station
Alexandria, Virginia 22314
- (6) Director, Naval Research Laboratory
Attention: Technical Information Division
Code 2627
Washington, D. C. 20375
- (6) Office of Naval Research
Attention: Code 102IP (ONRL DOC)
800 N. Quincy Street
Arlington, Virginia 22217
- (3) Office of Naval Research
Biophysics Program
Code 444
Arlington, Virginia 22217
- (1) Commanding Officer
Naval Medical Research and Development Command
National Naval Medical Center
Bethesda, Maryland 20014
- (1) Chief, Bureau of Medicine and Surgery
Department of the Navy
Washington, D. C. 20375
- (2) Technical Reference Library
Naval Medical Research Institute
National Naval Medical Center
Bethesda, Maryland 20014
- (1) Office of Naval Research Branch Office
Building 114, Section D
666 Summer Street
Boston, Massachusetts 02210
- (1) Office of Naval Research Branch Office
536 South Clark Street
Chicago, Illinois 60605

- (1) Office of Naval Research Branch Office
1030 East Green Street
Pasadena, California 91106
- (1) Commanding Officer
Naval Medical Research Unit No. 2
Box 14
APO San Francisco 96263
- (1) Commanding Officer
Naval Medical Research Unit No. 3
FPO New York 09527
- (1) Officer in Charge
Submarine Medical Research Laboratory
Naval Submarine Base, New London
Groton, Connecticut 06342
- (1) Scientific Library
Naval Aerospace Medical Research Institute
Naval Aerospace Medical Center
Pensacola, Florida 32512
- (1) Commanding Officer
Naval Air Development Center
Attn: Aerospace Medical Research Department
Warminster, Pennsylvania 18974
- (1) DIRECTOR
Naval Biosciences Laboratory
Building 844
Naval Supply Center
Oakland, California 94625
- (1) Commander, Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709
- (1) DIRECTORATE OF LIFE SCIENCES
Air Force Office of Scientific Research
Bolling Air Force Base
Washington, D. C. 20332
- (1) Commanding General
Army Medical Research and Development Command
Forrestal Building
Washington, D. C. 20314

- (1) Department of the Army
U. S. Army Science and
Technology Center - Far East
APO San Francisco 96328
- (1) Assistant Chief for Technology
Office of Naval Research, Code 200
800 N. Quincy Street
Arlington, Virginia 22217

